Lighting And Material Of Halo3

Hao Chen (haochen@bungie.com)

Bungie

Microsoft Research Asia

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRVM

Talk Overview

- Introduction
- Halo3 Lighting
- Halo3 Material Model
- HDR Rendering
- Results
- Acknowledgement
- Q&A

Halo2

per pixel light map

per vertex dominant lightdir+“ambientess”

Bump maps modulates light map color.

Objects sample LM and build fake lights.

Bumps only see single directional light

Bumps disappear In shadowy area

Specular from point lights only

LDR pipeline

Halo3

- What do we want?
  - global illumination
  - handle variety of lighting environments
  - consistent lighting everywhere
  - render bump maps “correctly”
  - complex material under complex lighting
  - HDR

Definitions

- Radiance
  - *L(ω)*: density of light energy through a given point in a given direction.
- Irradiance
  - *I:* incident radiance integrated over the hemisphere of the surface normal with the cosine lobe.
- BRDF
  - *f(V, L):* Bidirectional reflectance distribution function.
- Fresnel
  - *F:* Predicts ratio of reflected and transmitted light when light travels between different mediums.
  - *F0*: Reflectance at near normal incident angle.

BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM

Related Work

- Irradiance Volume [Greger98][ATI05]
- SH Irradiance Environment Map [Ramamoorthi01]
- Pre-computed Radiance Transfer [Sloan02]
- SH Light Maps [Good, Taylor05]
- Sky and Atmosphere [Preetham99][Hoffman02]
- Reflectance Models [CookTorrance82][Schlick94]
- Low Frequency Glossy Material [Kautz02] [Sloan03]
- Frequency Space Environment Map [Ramamoorthi02]

SH Irradiance Env Map



[Ramamoorthi00]

distant radiance in given direction

SH coefficients

diff solid angle

evaluated at given direction

SH Irradiance Env Map



- Irradiance distribution as SH vector.
- <3% error with just 9 terms [Basri Jacobs 01]
- Evaluate normal to get irradiance.
- Only represent a single point in space.
- Only for infinite lighting environment.
- What about local lighting?

Irradiance Volumes

[Greger 98]



[ATI 05]

- Spatially divide volume into cells.
- irradiance volume per cell.
- Interpolate between samples.
- Sharp shadow boundaries?
- Bump maps?

BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM

SH Light Map

Parameterize

GI Solver

linear

quadratic

Rendering



Compression

Store a SH Irradiance environment map per pixel over the surface

Other basis would work too, e.g. Half Life Basis, ZH, etc.

Sky and Sun

- HDR pipeline means starting with HDR light sources.
- Custom sky plugin for 3DStudio Max.
- Procedural Sky/Sun Model[Preetham99]
- CIE Sky Model
- Can also use HDR Cubemap



BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM

HDR Sky Example

NON FACETE NOBIS CALCITRARE VESTRVM

Parameterization



- UVAtlas (DirectX SDK)
  - Minimize distortion
  - Efficient Packing
  - Input “importance”
- Halo3 improvements:
  - small charts placement.
  - long and thin charts.
- > 80% texel utilization
  - Halo2: < 50%

Photon Mapping Farm



256 servers; 456 Processors; 1GB memory per proc;
see Luis's talk.

Compression



Signal Optimization:

- Compute signal gradient
- Resample to half dim.
- Preserve high freq charts
- Squeeze low freq charts

DXT Compression

- Use 2 DXT5 for each FP16
- Color space conversion
- 12 bits (2 DXT5 alpha) luminance
- 3:1 compression ratio

Details in a separate talk by Yaohua Hu.

NON FACETE NOBIS CALCITRARE VESTRVM

Rendering Passes



BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM

Object Lighting

sample surface

PRT

VS:
PRT/
Unshadowed

PS:
Unshadowed *
ratio

BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM

Optimization

- given a quadratic SH vecotor, i=0,...8
  - Pull out dominant light.
  - Store SH linear + dominant light.
  - In Shader, do N dot L + shirm(sh[] – c * Y(d), N)

See Peter Pike Sloan's talk "Stupid Spherical Harmonics Tricks".

Lighting Examples

NON FACETE NOBIS CALCITRARE VESTRVM

Lighting Examples



BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM

Lighting Examples

BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM

Lighting Examples

BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM

Lighting Examples



BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM

Lighting Examples



NON FACETE NOBIS CALCITRARE VESTRVM

Material Motivation



Halo3 Material Model

Diffuse

Area Specular



Environment Map

Analytical Specular

- The basic Idea
  - Separate material into diffuse parts and
  - Low, med, high freq glossy parts.
  - SH irradiance envmap for diffuse
  - New area specular model for low frequency glossy.
  - Prefiltered env map for mid frequency glossy.
  - BRDF evaluated directly with point lights for high freq.

NON FACETE NOBIS CALCITRARE VESTRVM

Cook Torrance BRDF

f

view light direction diffuse specular Fresnel specular lobe

[CookTorrance81]



[Ngan05]

D: microfacet distribution function

G: geomery term

BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM

Cook Torrance & Point Lights

- Option1: Evaluate directly in shader
  - E.g. [Dempski Viale 2005]
  - Somewhat expensive, not too bad.
- Option2: Store D, G, F, terms in textures.
- What about lights that are not point lights?
  - Need to integrate light from all directions.
  - Not trivial to do.

BUNGIE

Light Integration

??



SH irradiance env. map

BUNGIE

Light Integration in SH



Project light into SH basis.

Project BRDF and cosine term in SH basis

Dot product to convolve

Light Integration in SH Cont.

[Shilick85]



Pre-integration



Reflective symmetry means:

Isotropic BRDF = any coordinate frame values, and 8 V directions is enough.

C (i=0,2,3,6)



D (i=0,2,3,6)



C,D (i=7,8)

BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM

BUNGIE

BUNGIE

BUNGIE



BUNGIE

HDR Rendering

HDR Rendering

BUNGIE

Rendering Pipeline



BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM

Render Target Considerations

- Memory Size
- Render Speed
- Hardware Blend Support
- Dynamic Range
- Banding

Useable Exposure Range

BUNGIE

Xbox 360 Render Targets

	16f	10-bit 7e3	10-bit lin.	8-bit xRGB	16-bit lin.
Exposure Range	30 stops	3 stops	0 stops	0 stops	5 stops
Blend Support	NO	YES	YES	YES	YES
Memory Size	2x	1x	1x	1x	2x
Blend/Fill Rate	50%	50-100%	50-100%	[illegible]	50%



	8-bit xRGB + 8-bit xRGB
Exposure Range	7 stops
Blend Support	YES
Memory Size	2x
Blend/Fill Rate	25-50%

Detai
l
Bloom
(-7 stops)


Lack of sufficient dynamic range
makes the dark areas look hazy!
Artificially increase dynamic range...
BUNGIE


Bloom Intensity
0.0
1.0
2.0
HDR Intensity

Bloom Postprocessing

BUNGIE
NON FACETE NOBIS CALCITRARE VESTRVM


1.0
0.0
Linear LDR Valu
0.0
1.0
HDR Value
New
White Point
Clamped
Cubic Polynomial

No Tone Curve
BUNGIE

Tone Curve

BUNGIE

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRVM

NON FACETE NOBIS CALCITRARE VESTRVM

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRVM

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRVM

BUNGIE

NON FACETE NOBIS CALCITRARE VESTRVM

BUNGIE

Conclusion

- SH light map is a natural extension to the traditional light-mapping pipeline.
- Separating material into layers is a good approximation for all frequency reflectance.
- Area specular is critical for achieving seamless lighting and material integration.
- ALU is cheap, and will get cheaper, take maximum advantage of it.

Future Work

- Global Illumination with local, moving lights.
- GI for dynamic and semi-dynamic scenes.
- Better lighting basis (less ringing, higher frequency).
- Area specular model with complex transport.
- Measured BRDF.
- Non photo-realistic rendering.

Acknowledgement

- Authors
  - Bungie: Hao Chen, Ben Wallace, Chris Tchou, David Cook, Xi Wang, Yaohua Hu.
  - MSRA: Xinguo Liu, Zhipeng Hu, Xin Huang, Minmin Gong, Kun Zhou.
- Special Thanks
  - Bungie artists.
  - Peter Pike Sloan, Baining Guo, Harry Shum
  - Kutta Srinivasan, Matt Lee, Mikey Wetzel.

These slides will be available online
Feb. 25th, 2008
http://www.bungie.net/publications/

We're Hiring!
- Come see us at the Career Pavillion in Booth 913 (West Hall)

http://www.bungie.net/jobs/

NON FACETE NOBIS CALCITRARE VESTRVM

Q & A

BUNGIE